

# Producing High-Quality Plugs

How light quality, light quantity and carbon dioxide influence young plant production in greenhouses and indoor controlled environments.

Allison Hurt & Roberto Lopez

Every grower has their own protocol or recipe for producing plugs and rooted cuttings that works best with their growing environment, resources and needs. However, there's always room for improvement.

In this four-part series, we'll highlight our most recent greenhouse photoperiodic and supplemental lighting research utilizing light-emitting diodes (LEDs) during the production of both plugs and liners. Lastly, we'll introduce you to the exciting possibilities that exist with indoor vertical production of young plants utilizing sole-source LED lighting and carbon dioxide (CO<sub>2</sub>) injection.

Over the last decade, our research has quantified the influence of light quantity and quality during young plant production on numerous ornamental species. Our general recommendation for most young plants is to provide 70 to 90  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  of supplemental light to achieve a target daily light integral (DLI) of 10 to 12  $\text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$ . As a result, many growers are now producing young plants under supplemental lighting from high-pressure sodium (HPS) lamps, while others have made the investment in high-intensity LED fixtures (Figure 1A-D).

Most growers utilizing supplemental

lighting report that production time is often reduced and that young plants are of higher quality (increased rooting, stem diameter and branching) and often flower earlier upon transplant. Additionally, when the ambient DLI is low ( $<7 \text{ mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$ ) plugs of some species grown under LEDs providing  $\geq 10\%$  blue light are generally more compact (reduced leaf area and stem elongation), greener or have more pigmentation (anthocyanins) than those grown under HPS lamps or no supplemental lighting. Another group of growers use low-intensity LED photoperiodic lighting for 16 to 24  $\text{h}\cdot\text{d}^{-1}$  during young plant production and report reduced production time compared to no electric lighting (Figure 1E).

As the use of LED supplemental lighting increases, more questions surface. For example, since LEDs are much more energy-efficient to operate than HPS lamps, is there any benefit from running them continuously at a lower intensity even if the sun is out? Alternatively, is it more beneficial to run LEDs at a higher intensity in the morning, when it's cloudy and in the evening? There's value in addressing these questions, even though basic plant physiology tells us that plants can only utilize a certain quantity of light (light sat-

uration point) for photosynthesis and anything beyond this is wasted energy.

Additionally, most commercially available LED fixtures only provide red and blue light, which can make plant observation a little more challenging, especially when it's dark. Therefore, we also wanted to determine how the addition of white light, mainly for human applications, would affect plug production.

Our objectives were to quantify plug quality and production time under 1) continuous 16-hour or instantaneous threshold supplemental lighting with HPS lamps or high-intensity LED top lighting; and 2) under low-intensity LED photoperiodic lighting with and without far-red light and compare these methods to plugs receiving no electric lighting.

## The study

Seeds of *Petunia x hybrida* Ramblin' Peach Glow, *Impatiens walleriana* Accent Premium Salmon, *Gerbera jamesonii* Jaguar Deep Orange, wax begonia (*Begonia semperflorens* Bada Bing Scarlet) and tuberous begonia (*Begonia x tuberosa* Nonstop Rose Petticoat) were sown in 128-cell trays at a commercial greenhouse. One week later, trays were placed in a glass-glazed greenhouse at Michigan State University (MSU) in East Lansing, Michigan (lat. 40° N), with an air temperature set point of 72F (22C) and a vapor pressure deficit of 0.3 kPa maintained by injecting steam. Seedlings were hand irrigated as needed with reverse osmosis water supplemented with water-soluble fertilizer that provided 60 ppm nitrogen.

	Control	Supplemental lighting				Photoperiod lighting			
		HPS 16-h 70 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	HPS threshold 90 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	LED 16-h 70 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	LED threshold 90 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	Philips 16-h R:W:FR	Philips 16-h R:W	GE 16-h R:W:FR	GE 24-h R:W:FR
<i>Total no. of hours of operation</i>									
Nov.-Dec.	0	448	357	448	357	448	448	448	672
Jan.-Feb.	0	448	403	448	403	448	448	448	672
Feb.-Mar.	0	448	319	448	319	448	448	448	672
<i>DLI (<math>\text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}</math>)</i>									
Nov.-Dec.	4.8	9.7	9.4	9.9	10.0	5.7	5.4	5.5	5.7
Jan.-Feb.	5.5	9.9	10.1	9.7	10.3	5.9	5.8	5.6	6.0
Feb.-Mar.	7.5	12.9	11.8	11.7	11.2	8.5	8.6	8.7	7.9

Table 1. The total hours of operation and daily light integral (DLI) for no electric lighting, threshold and 16-hour high-pressure sodium (HPS) and light-emitting diode (LED) supplemental lighting, and 16- and 24-hour photoperiodic lighting for the four-week study.

Plug trays were placed under each of nine lighting treatments that included a control (natural daylength with no electric lighting), four supplemental lighting treatments with either HPS or LED fixtures, and four photoperiodic treatments with LED lamps (Figure 2). Supplemental lighting treatments consisted of high-intensity 200-watt LED fixtures (Philips GP-TOPlight DRW-MB) providing a light ratio (%) of 10:5:85 blue:green:red (B:G:R) or 400-watt HPS lamps (P.L. Light Systems) providing a PPFD of  $70 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  (on continuously for  $16\text{-h}\cdot\text{d}^{-1}$ ).

These same HPS lamps and LED fixtures were also used in instantaneous

threshold supplemental lighting treatments (on from 6:00 to 8:00 a.m. and 5:00 to 10:00 p.m. and only on between 8:00 a.m. to 5:00 p.m. when outside PPFD was  $<185 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  and switched off when  $>370 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) providing a PPFD of  $90 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ .


Photoperiodic lighting treatments consisted of screw-in low intensity 14-watt LED lamps providing  $2 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  of a light ratio (%) of 6:15:77:2 B:G:R:far-red (Philips GreenPower LED flowering DR:W) or screw-in low intensity 15-watt LED lamps providing 7:12:35:46 B:G:R:FR (DR:W:FR) for  $16\text{-h}\cdot\text{d}^{-1}$  or 10-W LEDs providing 3:17:48:32 B:G:R:FR (GE Arize

Greenhouse Pro Photoperiodic LED Lamp) for 16 or  $24\text{-h}\cdot\text{d}^{-1}$ .

Since our continuous supplemental lighting treatments ( $70 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) were on for  $16\text{-h}\cdot\text{d}^{-1}$  they operated a total of 448 or 672 hours during the four or six weeks of production, respectively (Table 1). In addition to sunlight, plugs under these treatments received an additional  $5 \text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$  of supplemental lighting.

On the other hand, our threshold supplemental lighting at  $90 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  provided  $2.3 \text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$  as fixtures were on each day from 6:00 to 8:00 a.m. and 5:00 to 10:00 p.m. On very cloudy days, the lamps could potentially continue operating from 8:00 a.m. to 5:00 p.m. and provide an additional  $4.2 \text{mol}\cdot\text{m}^{-2}\cdot\text{d}^{-1}$  of supplemental light. During January and February, our threshold lamps ran for 403 hours and provided a very similar DLI during the four weeks of production (Table 1).

After four or six weeks (begonia only), plugs were subsequently transplanted into 4- or 4.5-in. pots filled with a commercial soilless media and finished in a greenhouse with an air temperature set point of 68F (20C) (petunia, begonia and impatiens) under the LED top lighting fixtures mentioned above or 72F (22C) under HPS lamps (gerbera).

Stay tuned for our second article in the four-part series where we'll share the results of our study looking at the effects of light quality, quantity and duration on plug production. 

**Allison Hurt is a M.S. student and Roberto G. Lopez is an Assistant Professor and Controlled Environment/Floriculture Extension Specialist in the Department of Horticulture at Michigan State University. The authors gratefully acknowledge Ball Horticultural Company and Syngenta Flowers for providing seed, C. Raker and Sons for seed sowing, Philips Lighting, HortAmericas, The Western Michigan Greenhouse Association, and The Metro Detroit Flower Growers' Association for funding, and Nathan DuRussel for greenhouse assistance.**



Figure 1. Supplemental lighting of young plants utilizing A) high-pressure sodium lamps, B-D) various commercially available high-intensity light emitting diode (LED) fixtures or E) LED photoperiodic lighting.



Figure 2. A) Examples of high-pressure sodium (HPS), B) light-emitting diode (LED) Top lighting and C) photoperiod treatments (wire mesh was used to reduce light intensity).