

Prairie Fen Companion *Plant facts*

American burnweed, fireweed

Erechtites hieraciifolia



C=2

Color: Greenish-white

Parts indistinguishable

Plant type: Flowering perennial.

Bloom period: July – August.

Distinguishing characteristics: Leaves are alternate, with deeply or shallowly toothed elongate leaves. Greenish-white flowers $\frac{1}{2}$ to $\frac{3}{4}$ inch long bloom in clusters. Flowers are swollen at the base, with a brush-like tip, and may be mistaken for buds unless examined closely. The species then rapidly produces fluffy white seedheads. The stem is grooved and may be smooth or hairy; the plant has a weedy growth habit generally. Plants grow 1-8 feet tall.

Habitat quality: This species does not tolerate dense shade. It is fairly common after clearing of invasive shrubs, and is likely to establish first in areas that were burned, especially if brush piles were burned as part of clearing of invasives. It persists for the first few years after clearing. It is absent or uncommon in intact, high-quality prairie fen systems.

Where can I find this plant?



MICHIGAN STATE
UNIVERSITY




Developed by: Doug Landis and Anna Fiedler, MSU Department of Entomology. Funding support: National Fish and Wildlife Foundation, Lynn and Thelma MacCready Forest and Wildlife Endowment, MSU, and Hanes Trust of the Michigan Botanical Club. Partners: The Nature Conservancy, Michigan Natural Features Inventory, The Stewardship Network, Michigan DNR Landowner Incentive Program. For more information on native plants and prairie fens, go to www.nativeplants.msu.edu.

Where can I find this plant?




Indiana



New York



Ohio



Wisconsin

For more information on native plants and prairie fens, go to www.nativeplants.msu.edu

Key description

C=


Coefficient of conservatism. This is a value that ranges from 0 for non-native invasives to 10 for plants that would only be expected to be found in undisturbed, high quality plant communities. It is a general guideline for whether the plant would be likely to be found in an intact prairie fen (not filled in with shrubs and without invasive species). However, C values may be high for some species that are not found in prairie fen but would be part of another wetland such as a marsh. They are included here because they are a widely accepted measure of habitat quality in the Midwest (<http://1.usa.gov/FQAMethod>).

Flower type

Classifications here follow those of Newcomb's wildflower guide (<http://amzn.com/0316604429>).

Number of regular parts

The flower has this number of petals or petal-like parts that are symmetrical from the flower center (radial), with each similar to the other in shape, size, and color. There may be 3 to 7 regular parts. See image at right.



Aster

These flowers have regular parts and are symmetrical from the center, but there are more than 7. Asters have a set of disc flowers in the center of the flower and a set of ray flowers outside of the disc flowers, often called petals. They are one group (genus) within the family Asteraceae, and there are many species in this genus.

Flowers not readily obvious

The flowers of plants such as grasses, sedges, and cattail are not obvious and are often confused with the fruits (seeds) of these species. While this website does not include flower descriptions for these species, they do flower.

Irregular

The flower is not symmetrical from the center but is symmetrical down a line (bilateral). See image at right.



No flowers

A number of primitive plants, including ferns, do not flower but make spores in order to reproduce.

Parts indistinguishable

These species either have parts so small their number is difficult to determine or have no petal-like parts. This group includes goldenrods, other species with small individual flowers, and plants in the family Asteraceae that have more than 7 parts, but the parts do not form distinguishable, symmetrical ray flowers (which are often called petals).