



# CLIMATE CHANGE IN THE GREAT LAKES REGION


Don Scavia
University of Michigan









#### **Global Temperature**



## Global, Regional, Local








Global trends are more certain than regional trends

Natural variability plays a larger role at the regional scale

Local land use changes can alter severity of climate impacts

# What has Changed?



Changes are often discussed as averages ...

... but most environments are managed in terms of timing and extremes.



# **Today's Outline**

#### **Changing Temperatures**


- What has happened
- What is projected
- What are the likely impacts

#### **Changing Precipitation**

- What has happened
- What is projected
- What are the likely impacts



# What has Changed?




Changes are often discussed as averages ...

... but most environments are managed in terms of timing and extremes.



## **Observed Michigan Temperature**



Changes in Average Temperature (°F) from 1951-1980 to 1981-2010	
Annual	0.9
Winter	1.9
Spring	1.1
Summer	0.5
Fall	-0.1

Winter temps and overnight lows increased faster than annual averages.



#### **Observed Heat Waves**

Heat waves that pose risks to human health increased in most major Midwestern cities.

Increasing overnight, minimum temperatures increased faster, limiting relief during hot periods.

# Observed Change in Number of Harmful Heat Waves

Chicago, IL 1948–2011 (63 years)






**Detroit, MI** 1959–2011 (52 years)








#### **Migrating Plant Hardiness Zones**








Average extreme minimum temperatures migrated north.



## Longer Midwestern Growing Season



Growing season lengthened by ~1-2 weeks

Mostly due to earlier last winter frost in spring


Growing season in 2100 may be 1-2 months longer



## The Great Lakes are Warming

Average Great Lakes ice coverage declined 71% percent from 1973 to 2010

Wang et al., 2012



Lake Superior water is warming twice as fast as air


Lake Superior could have little to no open-lake ice cover during a typical winter within the next 30 years



## **Projection:** A Migrating Climate


Future generations will experience a fundamentally different climate.

By the end of this century,
Michigan summers will feel
like current summers in
Arkansas.






#### **Projected Midwest Temperature Increases**



#### Impacts in the Great Lakes Region

Changes in <u>temperature</u> and precipitation will impact both engineered and natural environments.







Fish
Water
Energy
Forests
Agriculture
Biodiversity
Public Health
Transportation








Birds and Wildlife
Tourism and Recreation




#### **More Hot Days Projected**



GLISA

#### **Impacts of Declining Lake Ice Cover**





Wang et al., 2012

- Fishing Industry: Harms whitefish spawning areas and increased wetland
- Coastal Zone: Loss of stable platform for recreation
- Navigation: Potentially lower water levels; lengthened shipping season



#### **Potential Impacts on Shipping**

#### **Every lost inch of water depth:**

- Reduces cargo capacity 50-270 tons
- Costs \$10k-30k per transit






...but less lake ice cover also allows for a longer shipping season




# **Lake Level Projections**





#### **Projected Shifts in Forest Types**



#### Impacts on Biodiversity

- Amplified existing stressors, including sensitivity to land and water use
- Some species will need to migrate to keep up with the pace of warming ...

... but, large agricultural areas and the Great Lakes are major obstacles to migration







#### **Impacts on Agriculture**



- Some crops may benefit in the near future from increasing carbon dioxide concentrations until negated by warmer temperatures.
- Perennial crops may be more vulnerable to the pace of climate change and may face greater adaptation challenges.



#### Agriculture Vulnerabilities: Spring 2012 Cherry Crops


- The early warming was extreme weather event
- The seasonal warming fits a pattern of a more variable climate



- The early warming followed by a normal hard freeze was devastating to cherry buds
- \$92 million loss from tart cherries alone



# What has Changed?




Changes are often discussed as averages ...

... but most environments are managed in terms of timing and extremes.




## **Observed Michigan Precipitation**



Changes in Total Precipitation (%) from 1951-1980 to 1981-2010	
Annual	5.0
Winter	5.6
Spring	1.3
Summer	-1.0
Fall	15.8

Precipitation increased in Winter and Fall but remained stable or have declined during Spring and Summer.


#### **Observed Extreme Precipitation Intensity**



1% of precipitation events increased from 1958 to 2007.




#### **Observed Extreme Precipitation Frequency**



Frequency of heaviest 1% of precipitation events increased from 1958 to 2007.



# **Observed Snowfall Change**



Snowfall increased in the North & decreased in the South



#### **Changing Winter Precipitation**





- Lake effect snow increased in some areas.
- But, shorter winters led to more falling as rain.



Warmer surface temperatures also reduced snow accumulation.



## **Projected Precipitation**




Percent Change

-15 - -10-10 --5


5-0 0-5 5-10

15 - 20

Winter +5 to 20%



(2041-2070 minus 1971-2000)



NARCCAP, Precipitation Change, Fall (2041-2070 minus 1971-2000)



Annual +5 to 15%

**Spring** +0 to 20%

Summer +10 to -10%



Kunkel (2011)

GLISA

#### **Impacts in the Great Lakes Region**

Changes in temperature and <u>precipitation</u> will impact both engineered and natural environments.







Fish
Water
Energy
Forests
Agriculture
Biodiversity
Public Health
Transportation







Birds and Wildlife
Tourism and Recreation



# Flooding and Stormwater

Stronger and more frequent extreme events amplify flood risks.













#### **Impacts on Agriculture**


- Increasing intensity of severe storms increases the risk of runoff and erosion.
- Shifts in the timing of precipitation will affect field preparation time in spring.




- Some crops may benefit in the near future from increasing carbon dioxide concentrations until negated by warmer temperatures.
- Perennial crops may be more vulnerable to the pace of climate change and may face greater adaptation challenges.




## **Conspiring Changes: Water Quality**



## **Algal Blooms and Fish Kills**



## How will we adapt?



## Are we preparing to adapt?









# Extra Slides

# **Projected Snowfall Days**

